

EMBARGOED FOR RELEASE until May 19, 2016 – 5:00 PM (ET)

Contacts: Tracy Hampton • (312) 339-9067 • thampton@nasw.org
Bob Henkel • (202) 557-8360 • bhenkel@asn-online.org

SIMPLE PUBLIC HEALTH INTERVENTION MAY PREVENT CHRONIC KIDNEY DISEASE

Program may be especially helpful in developing countries

Highlights

- Kidney function remained unchanged among hypertensive adults in communities assigned to a public health intervention for general practitioners and community health workers, whereas kidney function significantly declined among those who received usual care.
- Individuals in the communities with the intervention were half as likely as other individuals to experience a >20% decline in kidney function within 5 years after the intervention was stopped.

An estimated 10% to 15% of adults in the world have chronic kidney disease.

Washington, DC (May 19, 2016) — A simple and inexpensive public health intervention helped prevent many cases of chronic kidney disease (CKD) in Pakistan. The intervention, which is described in a study in the *Clinical Journal of the American Society of Nephrology* (CJASN), will be especially helpful for protecting the kidney health of people living in developing countries.

Up to 500 million people in the world have CKD, which has become one of the most rapidly escalating causes of death globally. Patients with advanced CKD require dialysis or transplantation to sustain life, but these therapies are unaffordable for the vast majority of people with CKD in developing nations. Unfortunately, there is dearth of information on public health interventions to preserve kidney health and prevent CKD, even though certain lifestyle factors and conditions (such as hypertension and diabetes) are known to increase the risk of developing the disease.

To address this, Tazeen Jafar, MD, MPH (Duke NUS Medical School, in Singapore, Duke Global Health Institute, in Durham, NC) and her colleagues from Pakistan, Singapore, and the United Kingdom assessed the effects of a combined public health intervention on the kidney health of hypertensive adults in Pakistan's general population. The study based at the Aga Khan University, Karachi, included 1271 individuals, and the intervention was delivered over 2 years. It included training of community health workers on aspects of a healthy lifestyle (such as improving diet, stopping smoking, increasing

physical activity, and taking prescribed blood pressure–lowering medications) and training of community general practitioners on the latest standards related to managing hypertension.

After 7 years of follow up—5 years after cessation of the intervention—kidney function remained unchanged among adults in the communities assigned to the combined intervention, whereas kidney function significantly declined among those who received usual care. Individuals in the communities with the combined intervention were half as likely as other individuals to experience a >20% decline in kidney function.

“We show that such a practical model based on primary care doctor training coupled with life style advice from non-physician health workers is likely to have a long-term benefit on preserving kidney function at a population level,” said Dr. Jafar. “These simple strategies can be implementable in other low- and middle-income countries with similar risk factor burden and health systems infrastructure.”

In an accompanying editorial, Min Jun MScMed, PhD and Brenda Hemmelgarn, MD, PhD (University of Calgary, in Canada) noted that adequate blood pressure control among high-risk patient groups such as those with CKD is as low as 13.2%, and awareness of CKD is lower in developing countries than in developed countries. “It follows that simple interventions based on education and communication of the importance of established prevention strategies including blood pressure management may have a significant impact at the population level,” they wrote. “This therefore warrants further consideration including the assessment of the cost-effectiveness and sustainability of prevention strategies specific to CKD management in developing countries.”

Study co-authors include John Allen, PhD; Imtiaz Jehan, MD, MSc; Aamir Hameed, MD; Seyed Ehsan Saffari, PhD; Shah Ebrahim, MD; Neil Poulter, F Med Sci; and Nish Chaturvedi, MD.

Disclosures: The authors reported no financial disclosures.

The article, entitled “Home Health Education and General Practitioner Training in Hypertension Management: Long Term Effects on Kidney Function,” will appear online at <http://cjasn.asnjournals.org/> on May 19, 2016, doi: 10.2215/CJN.05300515.

The editorial, entitled “Strategies for blood pressure control in developing countries and effects on kidney function,” will appear online at <http://cjasn.asnjournals.org/> on May 19, 2016.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions,

drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Founded in 1966, and with nearly 16,000 members, the American Society of Nephrology (ASN) leads the fight against kidney disease by educating health professionals, sharing new knowledge, advancing research, and advocating the highest quality care for patients.

#

Tweet: Simple public health intervention may prevent chronic kidney disease.
<http://www.bit.ly/ASN-XXXX>. Study Twitter handle: @COBRASStudy. An extension of COBRA in rural Bangladesh, Pakistan, and Sri Lanka funded jointly by the Wellcome Trust/MRC and DfID, UK has a dedicated Facebook page:
<https://www.facebook.com/cobra.bps/>

Facebook: A simple and inexpensive urine test routinely done in family doctors' offices can identify people who are silently undergoing rapid kidney function decline, says a study in the *Journal of the American Society of Nephrology*. The test could lead to earlier and more effective treatments, lowering risks of kidney failure and death. Some 60 million people globally have chronic kidney disease, but many don't know it, as they have no symptoms until later stages of disease.

Media:

Dharshini Subbiah - dharsini.subbiah@duke-nus.edu.sg