



PRESS RELEASE

ASN Contacts:

Christine Feheley (202) 640-4638 | cfeheley@asn-online.org

Tracy Hampton thampton@nasw.org

RACIAL/ETHNIC DISPARITIES IN STROKE-PREVENTION AMONG PATIENTS UNDERGOING DIALYSIS

Study reveals that minorities are less likely to fill prescriptions for blood thinners

Highlights

- Among patients with kidney failure and atrial fibrillation, Black, Hispanic White, and Asian patients filled prescriptions of stroke-preventive medications less often than non-Hispanic White patients, and they were more likely to experience stroke.
- Equalizing the distribution of these medications would prevent 7%–12% of the stroke disparity among racial/ethnic minorities.

Washington, DC (February 20, 2020) — In a study of patients with kidney failure and atrial fibrillation, racial/ethnic minorities experienced higher rates of stroke compared with non-Hispanic White patients, and they were less likely to fill prescriptions of stroke-preventive medications. The findings, which appear in an upcoming issue of *JASN*, indicate that equalizing the distribution of such prescriptions may help address stroke-related disparities among patients.

Because patients with kidney failure and atrial fibrillation, or an irregular heartbeat, are at risk of developing strokes, they may benefit from taking blood thinners as a preventive measure. Prior studies of patients with atrial fibrillation, with or without kidney failure, have reported that minority racial/ethnic groups face a higher risk of stroke compared with Whites.

To investigate the potential causes of such disparities, a team led by Paul L. Kimmel, MD (National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health) examined information from a national registry of patients with kidney failure.

For the study, the researchers analyzed information from the United States Renal Data System to identify patients with kidney failure who initiated hemodialysis from 2006 to 2013, and then they identified those with a subsequent atrial fibrillation diagnosis and Medicare Part A/B/D insurance coverage.

Among 56,587 patients who were followed for one year, the number of strokes per 1,000 people were 84, 94, 97, and 102 in non-Hispanic White, Black, Hispanic White, and Asian patients, respectively.

Black, Hispanic White, and Asian patients filled prescriptions of the blood-thinner warfarin less often than non-Hispanic White patients, and they were more likely to experience stroke. Specifically, Black, Hispanic White, and Asian patients were 10%, 17%, and 28% less likely than non-Hispanic Whites to fill a warfarin prescription, respectively, and they were 13%, 15%, and 16% more likely to experience stroke. (The database did not provide information on whether patients were prescribed drugs, only whether they filled prescriptions.)

Analyses suggested that equalizing the warfarin distribution to that in the non-Hispanic White patient population would prevent 7%, 10%, and 12% of the stroke disparity among Black, Hispanic White, and Asian patients, respectively.

“Institutional features and medical choices may play key roles underlying differences between prescriptions received by members of different groups. Our estimates suggest equalization of prescription of warfarin, a relatively inexpensive anticoagulant medication, across all ethnic/racial groups would be associated with decreased stroke rates in Black, Asian, and Hispanic White patients on hemodialysis,” said Dr. Kimmel.

Study co-authors include Salina P. Waddy, MD; Allen J. Solomon, MD; Adan Z. Becerra, PhD; Julia B. Ward, PhD, MPH; Kevin E. Chan, MD, MSc; Chyng-Wen Fwu, DrPH; Jenna M. Norton, MPH; Paul W. Eggers, PhD; and Kevin C. Abbott, MD, MPH.

Disclosures: Adan Z. Becerra, Julia B. Ward, and Chyng-Wen Fwu were supported by the National Institute of Diabetes and Digestive and Kidney Diseases contract HHSN275201700074.

The article, entitled “Racial/Ethnic Disparities in Atrial Fibrillation Treatment and Outcomes among Dialysis Patients in the United States,” will appear online at <http://jasn.asnjournals.org/> on February 20, 2020, doi: 10.1681/ASN.2019050543.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not

ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Since 1966, the American Society of Nephrology (ASN) has been leading the fight to prevent, treat, and cure kidney diseases throughout the world by educating health professionals and scientists, advancing research and innovation, communicating new knowledge, and advocating for the highest quality care for patients. ASN has more than 21,000 members representing 131 countries. For more information, visit www.asn-online.org.

The content of this news release is solely the responsibility of the authors and does not necessarily represent the official views or imply endorsement of the National Institutes of Health.

#

Tweet: Racial/ethnic disparities in stroke-prevention among patients undergoing dialysis.
Twitter: @niddkgov

Facebook: In a study of patients with kidney failure and atrial fibrillation, racial/ethnic minorities experienced higher rates of stroke compared with White patients, and they were less likely to receive prescriptions of stroke-preventive medications. The findings, which appear in *JASN*, indicate that equalizing the distribution of such prescriptions may help address stroke-related disparities among patients.